

Solar wind data assimilation in an operational context

Use of near-real-time data and the forecast value of an L5 monitor


Harriet Turner


Supervisors: Mathew Owens, Matthew Lang, Mike Marsh and
Siegfried Gonzi

THE SOLAR WIND

Constant stream of charged particles that flows off the Sun and fills the heliosphere (solar system)

Comprised mostly of electrons, protons and ions






*Not to scale

SOLAR WIND STRUCTURE


Solar wind drags out the Sun's magnetic field

Pulled into an Archimedean spiral due to the Sun's rotation

Solar wind flow is **mostly radial**




AS A TIME SERIES



CORONAL MASS EJECTIONS

Coronal mass ejections (CMEs) are huge eruptions of solar material and are the main driver of severe space weather

They propagate through the solar wind, so the background conditions affect their speed and arrival time





2012/07/06 23:12

WHY SHOULD WE CARE?

Space weather (the changing plasma conditions in near-Earth space) poses a significant threat to modern technology

Extreme space weather is in the UK's National Risk Register

Impact (of the reasonable worst case scenario using the impact indicators below)	Level E		7 25†		
	Level D	34*	12 13 29		
	Level C	18 28 33* 36*	14 19 21 26† 27* 38	2 3 6* 15 16 17 20	
	Level B	30	24	35*	4 5 9* 10* 11* 23 32* 37
	Level A		8* 22	31	1
		< 1 in 500	1 to 5 in 500	5 to 25 in 500	25 to 125 in 500
Likelihood (of the reasonable worst case scenario of the risk occurring in the next year)					

*Risk not plotted in the 2017 NRR | †COVID-19 is not included in the risk matrix and is therefore not included in these risks

Malicious Attacks

1. Attacks on publicly accessible locations
2. Attacks on infrastructure
3. Attacks on transport
4. Cyber attacks
5. Smaller scale CBRN attacks
6. Medium scale CBRN attacks
7. Larger scale CBRN attacks
8. Undermining the democratic process*

Serious and Organised Crime

9. Serious and organised crime – vulnerabilities*
10. Serious and organised crime – prosperity*
11. Serious and organised crime – commodities*

Environmental Hazards

12. Coastal flooding
13. River flooding
14. Surface water flooding
15. Storms
16. Low temperatures
17. Heatwaves
18. Droughts
19. Severe space weather
20. Volcanic eruptions
21. Poor air quality
22. Earthquakes
23. Environmental disasters overseas
24. Wildfires

Human and Animal Health

25. Pandemics†
26. High consequence infectious disease outbreaks†
27. Antimicrobial resistance*
28. Animal diseases

Major Accidents

29. Widespread electricity failures
30. Major transport accidents
31. System failures
32. Commercial failures*
33. Systematic financial crisis*
34. Industrial accidents – nuclear*
35. Industrial accidents - non nuclear*
36. Major fires*

Societal Risks

37. Industrial action
38. Widespread public disorder

Impact (of the reasonable worst case scenario using the impact indicators below)	Level E			7 25† Pandemics		
	Level D	34*		12 13 29		
	Level C		18 28 33* 36*	14 19 21 26† 27* 38	2 3 6* 15 16 17 20	
	Level B	30	24	35*	4 5 9* 10* 11* 23 32* 37	1
	Level A			8* 22	31	
		< 1 in 500	1 to 5 in 500	5 to 25 in 500	25 to 125 in 500	> 125 in 500
Likelihood						

*Risk not plotted in the 2017 NRR | [†]COVID-19 is not included in the risk matrix and is therefore not included in these risks

Malicious Attacks

- Attacks on publicly accessible locations
- Attacks on infrastructure
- Attacks on transport
- Cyber attacks
- Smaller scale CBRN attacks
- Medium scale CBRN attacks
- Larger scale CBRN attacks
- Undermining the democratic process*

Serious and Organised Crime

- Serious and organised crime – vulnerabilities*
- Serious and organised crime – prosperity*
- Serious and organised crime – commodities*

Environmental Hazards

- Coastal flooding
- River flooding
- Surface water flooding
- Storms
- Low temperatures
- Heatwaves
- Droughts
- Severe space weather
- Volcanic eruptions
- Poor air quality
- Earthquakes
- Environmental disasters overseas
- Wildfires

Human and Animal Health

- Pandemics[†]
- High consequence infectious disease outbreaks[†]
- Antimicrobial resistance*
- Animal diseases

Major Accidents

- Widespread electricity failures
- Major transport accidents
- System failures
- Commercial failures*
- Systematic financial crisis*
- Industrial accidents – nuclear*
- Industrial accidents - non nuclear*
- Major fires*

Societal Risks

- Industrial action
- Widespread public disorder

Impact (of the reasonable worst case scenario using the impact indicators below)	Level E			7 25† Pandemics		
	Level D	34*		12 13 29		
	Level C		18 28 33* 36*	14 19 21 26† 27* 38	2 3 6* 15 16 17 20	
	Level B	30	24	35*	4 5 9* 10* 11* 23 32* 37	1
	Level A			8* 22	31	
		< 1 in 500	1 to 5 in 500	5 to 25 in 500	25 to 125 in 500	> 125 in 500
Likelihood						

*Risk not plotted in the 2017 NRI | [†]COVID-19 is not included in the risk matrix and is therefore not included in these risks

Malicious Attacks

1. Attacks on publicly accessible locations
2. Attacks on infrastructure
3. Attacks on transport
4. Cyber attacks
5. Smaller scale CBRN attacks
6. Medium scale CBRN attacks
7. Larger scale CBRN attacks
8. Undermining the democratic process*

Serious and Organised Crime

9. Serious and organised crime – vulnerabilities*
10. Serious and organised crime – prosperity*
11. Serious and organised crime – commodities*

Environmental Hazards

12. Coastal flooding
13. River flooding
14. Surface water flooding
15. Storms
16. Low temperatures
17. Heatwaves
18. Droughts
19. Severe space weather
20. Volcanic eruptions
21. Poor air quality
22. Earthquakes
23. Environmental disasters overseas
24. Wildfires

Human and Animal Health


25. Pandemics[†]
26. High consequence infectious disease outbreaks[†]
27. Antimicrobial resistance*
28. Animal diseases

Major Accidents

29. Widespread electricity failures
30. Major transport accidents
31. System failures
32. Commercial failures*
33. Systematic financial crisis*
34. Industrial accidents – nuclear*
35. Industrial accidents - non nuclear*
36. Major fires*

Societal Risks

37. Industrial action
38. Widespread public disorder



Malicious Attacks

1. Attacks on publicly accessible locations
2. Attacks on infrastructure
3. Attacks on transport
4. Cyber attacks
5. Smaller scale CBRN attacks
6. Medium scale CBRN attacks
7. Larger scale CBRN attacks
8. Undermining the democratic process*

Serious and Organised Crime

9. Serious and organised crime – vulnerabilities*
10. Serious and organised crime – prosperity*
11. Serious and organised crime – commodities*

Environmental Hazards

12. Coastal flooding
13. River flooding
14. Surface water flooding
15. Storms
16. Low temperatures
17. Heatwaves
18. Droughts
19. Severe space weather
20. Volcanic eruptions
21. Poor air quality
22. Earthquakes
23. Environmental disasters overseas
24. Wildfires

Human and Animal Health

25. Pandemics†
26. High consequence infectious disease outbreaks†
27. Antimicrobial resistance*
28. Animal diseases

Major Accidents

29. Widespread electricity failures
30. Major transport accidents
31. System failures
32. Commercial failures*
33. Systematic financial crisis*
34. Industrial accidents – nuclear*
35. Industrial accidents - non nuclear*
36. Major fires*

Societal Risks

37. Industrial action
38. Widespread public disorder

Impact (of the reasonable worst case scenario using the impact indicators below)	Level E			7 25†		
	Level D	34*		12 13 29		
	Level C		18 28 33* 36*	14 19 21 26† 27* 38	2 3 6* 15 16 17 20	Heatwaves
	Level B	30	24	35*	4 5 9* 10* 11* 23 32* 37	1
	Level A			8* 22	31	
		< 1 in 500	1 to 5 in 500	5 to 25 in 500	25 to 125 in 500	> 125 in 500
Likelihood						

*Risk not plotted in the 2017 NRR | †COVID-19 is not included in the risk matrix and is therefore not included in these risks

Malicious Attacks

1. Attacks on publicly accessible locations
2. Attacks on infrastructure
3. Attacks on transport
4. Cyber attacks
5. Smaller scale CBRN attacks
6. Medium scale CBRN attacks
7. Larger scale CBRN attacks
8. Undermining the democratic process*

Serious and Organised Crime

9. Serious and organised crime – vulnerabilities*
10. Serious and organised crime – prosperity*
11. Serious and organised crime – commodities*

Environmental Hazards

12. Coastal flooding
13. River flooding
14. Surface water flooding
15. Storms
16. Low temperatures
17. Heatwaves
18. Droughts
19. Severe space weather
20. Volcanic eruptions
21. Poor air quality
22. Earthquakes
23. Environmental disasters overseas
24. Wildfires

Human and Animal Health

25. Pandemics†
26. High consequence infectious disease outbreaks†
27. Antimicrobial resistance*
28. Animal diseases

Major Accidents

29. Widespread electricity failures
30. Major transport accidents
31. System failures
32. Commercial failures*
33. Systematic financial crisis*
34. Industrial accidents – nuclear*
35. Industrial accidents - non nuclear*
36. Major fires*

Societal Risks

37. Industrial action
38. Widespread public disorder

Impact (of the reasonable worst case scenario using the impact indicators below)	Level E			7 25†		
	Level D	34*		12 13 29		
	Level C		18 28 33* 36*	14 19 21 26† 27* 38	20	
	Level B	30	24	35*	4 5 9* 10* 11* 23 32* 37	1
	Level A			8* 22	31	
		< 1 in 500	1 to 5 in 500	5 to 25 in 500	25 to 125 in 500	> 125 in 500
Likelihood						

Severe space weather

Malicious Attacks

1. Attacks on publicly accessible locations
2. Attacks on infrastructure
3. Attacks on transport
4. Cyber attacks
5. Smaller scale CBRN attacks
6. Medium scale CBRN attacks
7. Larger scale CBRN attacks
8. Undermining the democratic process*

Serious and Organised Crime

9. Serious and organised crime – vulnerabilities*
10. Serious and organised crime – prosperity*
11. Serious and organised crime – commodities*

Environmental Hazards

12. Coastal flooding
13. River flooding
14. Surface water flooding
15. Storms
16. Low temperatures
17. Heatwaves
18. Droughts
19. Severe space weather
20. Volcanic eruptions
21. Poor air quality
22. Earthquakes
23. Environmental disasters overseas
24. Wildfires

Human and Animal Health

25. Pandemics†
26. High consequence infectious disease outbreaks†
27. Antimicrobial resistance*
28. Animal diseases


Major Accidents

29. Widespread electricity failures
30. Major transport accidents
31. System failures
32. Commercial failures*
33. Systematic financial crisis*
34. Industrial accidents – nuclear*
35. Industrial accidents - non nuclear*
36. Major fires*

Societal Risks

37. Industrial action
38. Widespread public disorder

*Risk not plotted in the 2017 NRR | †COVID-19 is not included in the risk matrix and is therefore not included in these risks




CURRENT FORECASTING

Typically a **coronal model** coupled to a **heliospheric model**




CURRENT FORECASTING

Typically a **coronal model** coupled to a heliospheric model




CURRENT FORECASTING

Typically a **coronal model** coupled to a heliospheric model




CURRENT FORECASTING

Typically a **coronal model** coupled to a heliospheric model




CURRENT FORECASTING

Typically a **coronal model** coupled to a heliospheric model




CURRENT FORECASTING

Typically a **coronal model** coupled to a heliospheric model




CURRENT FORECASTING

Typically a **coronal model** coupled to a heliospheric model




CURRENT FORECASTING

Typically a **coronal model** coupled to a heliospheric model




CURRENT FORECASTING

Typically a coronal model coupled to a **heliospheric model**




CURRENT FORECASTING

Typically a coronal model coupled to a **heliospheric model**




CURRENT FORECASTING

Typically a coronal model coupled to a **heliospheric model**



CURRENT FORECASTING

Typically a coronal model coupled to a **heliospheric model**



DATA ASSIMILATION (DA)

Data assimilation combines prior information, usually from a model, with observations to find an optimum estimation of reality


Used in numerical weather prediction and has led to large forecast improvements

Underused in space weather

DA IN SPACE WEATHER

DA is in its infancy in space weather forecasting

Has been used in 3 main areas – the photosphere, solar wind and ionosphere



Photosphere –
lowest layer of the
Sun's atmosphere
that is observable

Solar wind


Ionosphere –
where Earth's atmosphere
reaches space


DA IN THE SOLAR WIND


I have been using the Burger Radius Variational Data Assimilation (BRaVDA) scheme developed at the University of Reading (Lang et al., 2019)

Combines observations with a steady-state solar wind model

Prior inner boundary condition from a coronal model









Time = t_0

✗ = observation




Time = t_0

✗ = observation




Time = $t_0 - \Delta t$

✗ = observation




Time = $t_0 - \Delta t$

✗ = observation




Time = $t_0 - \Delta t$

✗ = observation




Time = $t_0 - \Delta t$

✗ = observation




Time = $t_0 - \Delta t$

✗ = observation




Time = $t_0 - \Delta t$

✗ = observation




Time = $t_0 - \Delta t$

✗ = observation




Time = $t_0 - \Delta t$

✗ = observation




Time = t_0

✗ = observation




Time = t_0


✗ = observation



Time = t_0

✗ = observation






Time = $t_0 + \Delta t$

✗ = observation


CURRENT FORECASTING

Typically a **coronal model** coupled to a **heliospheric model**




CURRENT DA IMPLEMENTATION

Typically a **coronal model** coupled to a **heliospheric model**




CURRENT DA IMPLEMENTATION

Typically a **coronal model** coupled to a **heliospheric model**




CURRENT DA IMPLEMENTATION

Typically a coronal model coupled to a heliospheric model



CURRENT DA IMPLEMENTATION

Typically a coronal model coupled to a heliospheric model




DIVERGENCE OF METHODS

Previous work (Lang et al., 2021) have used the reduced-physics model “HUXt” (Owens et al., 2020) for analysis

For simplicity, I have used corotation – essentially assuming the Sun remains constant as it rotates


HUXt METHOD

Typically a **coronal model** coupled to a **heliospheric model**




HUXt METHOD

Typically a **coronal model** coupled to a **heliospheric model**




HUXt METHOD

Typically a **coronal model** coupled to a **heliospheric model**




HUXt METHOD

Typically a **coronal model** coupled to a **heliospheric model**




HUXt METHOD

Typically a **coronal model** coupled to a **heliospheric model**




COROTATION METHOD

Typically a coronal model coupled to a heliospheric model




COROTATION METHOD

Typically a **coronal model** coupled to a **heliospheric model**



COROTATION METHOD

Typically a **coronal model** coupled to a **heliospheric model**



OBSERVATIONS – STEREO


Solar Terrestrial Relations Observatory Ahead and Behind –
STEREO-A and B

Earth-like orbits, but separating in longitude

Launched in 2007

Communication with STEREO-B lost in 2014

NASA's STEREO (with SDO) Sees the Entire Sun



The two **STEREO** spacecraft reach equidistant positions between themselves and Earth on Sept. 1, 2012.


Drawing gives the relative orbital positions of both STEREO spacecraft for each year from June 2007 to June 2015.
(Not to scale)


OBSERVATIONS – ACE & DSCOVR

Advanced Composition Explorer – ACE (launched 1997)

Deep Space Climate Observatory – DSCOVR (launched 2015)

Both give near-Earth observations from L1









*Not to scale

DA OUTPUT – POLAR


CR2096




DA OUTPUT – TIME SERIES




FORECASTING USING OUTPUT




FORECASTING USING OUTPUT



FORECASTING USING OUTPUT



FORECASTING USING OUTPUT




OPERATIONAL SOLAR WIND DA

For BRaVDA to be operational, it needs to work with real time data


Previous experiments have used science level data, which has been pre-processed to provide a “cleaner” dataset (e.g. Lang et al., 2021 and Turner et al., 2022)

- Removing data gaps, erroneous observations and biases

REAL TIME ISSUES



REAL TIME ISSUES




BRaVDA EXPERIMENTS

Assimilating all sources of observations – STEREO-A, B and ACE


Evaluate the forecast accuracy at the three observation locations – verification time series is the science data

Comparing the prior, real time and science forecasts


BRaVDA EXPERIMENTS




TAYLOR DIAGRAM




TAYLOR DIAGRAM




TAYLOR DIAGRAM



TAYLOR DIAGRAM



TAYLOR DIAGRAM



FUTURE SOLAR WIND DA

The ESA Vigil mission is (hopefully) going to be launched in 2027 (ish) to the L5 point

“The mission will give us advanced warning of oncoming solar storms and therefore more time to protect spacecraft in orbit, infrastructure on the ground and explorers now and in the future, unshielded by Earth’s magnetic field and vulnerable to our star’s violent outbursts.” – ESA, 2023

Could be useful for future operational solar wind DA

L5 & L1 SIMULATION


Combinations of spacecraft separated by approximately 60° in longitude

Historic data to see how well a potential L5 and L1 pairing could work for solar wind DA


4 time periods

L5 & L1 SIMULATION


02/05/2008 - 30/08/2008







STEREO-A: 27/05/2009 - 06/05/2010
STEREO-B: 30/07/2009 - 22/01/2010



25/10/2013 - 09/02/2014




▲ STEREO-A ■ STEREO-B ● Earth




— Prior (no DA) — L1 only (science DA) - - L1 only (NRT DA) — L1 & L5 (science DA) - - L1 & L5 (NRT DA)


L5 & L1 AVERAGE




L5 & L1 AVERAGE




L5 & L1 AVERAGE



L5 & L1 AVERAGE



L5 & L1 AVERAGE



FUTURE WORK

I am currently working on a semi-operational version of BRaVDA

All the pieces are there but need pulling together

Hoping for a daily updated forecast – a world's first using solar wind DA

Something about a thesis?

CONCLUSIONS

Knowledge of the solar wind is important for timely and accurate space weather forecasting

The BRaVDA scheme reconstructs a full solar rotation using available observations

Leads to improved solar wind forecasts

A future L5 & L1 spacecraft pairing could be useful for operational solar wind DA

REFERENCES

ESA, 2023. URL: https://www.esa.int/Space_Safety/Vigil

Lang, M., & Owens, M. J. (2019). A Variational Approach to Data Assimilation in the Solar Wind. *Space Weather*, 17(1), 59 – 83. DOI: 10.1029/2018SW001857.

Lang, M., Witherington, J., Owens, M. J., & Turner, H. (2021) Improving solar wind forecasting using data assimilation. *Space Weather*, 1 – 23.

National Risk Register. URL: <https://www.gov.uk/government/publications/national-risk-register-2020>

Owens, M. J., et al. A Computationally Efficient, Time-Dependent Model of the Solar Wind for use as a Surrogate to Three-Dimensional Numerical Magnetohydrodynamic Simulations. *Solar Physics*, 295:43. DOI: 10.1007/s11207-020-01605-3.

Turner, H., Owens, M. J., Lang, M., Gonzi, S., & Riley, P. (2022). Quantifying the effect of ICME removal and observation age for in situ solar wind data assimilation. *Space Weather*, 20. DOI: 10.1029/2022SW003109.

Paper

Turner, H., Lang, M., Owens, M., Smith, A., Riley, P., Marsh, M., & Gonzi, S. (2023). Solar wind data assimilation in an operational context: use of near-real-time data and the forecast value of an L5 monitor. *Space Weather*, 21, e2023SW003457. <https://doi.org/10.1029/2023SW003457>

